


BACLOFEN- baclofen tablet
TruPharma LLC

BACLOFEN TABLETS, USP

DESCRIPTION

Baclofen, USP is a muscle relaxant and antispastic.

Its chemical name is 4-amino-3-(4-chlorophenyl)-butanoic acid. The structural formula is:



C₁₀H₁₂ClNO₂ M.W. 213.66

Baclofen, USP is a white to off-white odorless or practically odorless crystalline powder. It is slightly soluble in water, very slightly soluble in methanol and insoluble in chloroform.

Each tablet, for oral administration, contains 5 mg, 10 mg or 20 mg baclofen, USP. In addition, each tablet contains the following inactive ingredients: Microcrystalline Cellulose, Pregelatinized Starch, Sodium Starch Glycolate, Colloidal Silicon dioxide and Magnesium Stearate.

CLINICAL PHARMACOLOGY

The precise mechanism of action of baclofen is not fully known. Baclofen is capable of inhibiting both monosynaptic and polysynaptic reflexes at the spinal level, possibly by hyperpolarization of afferent terminals, although actions at supraspinal sites may also occur and contribute to its clinical effect. Although baclofen is an analog of the putative inhibitory neurotransmitter gamma-aminobutyric acid (GABA), there is no conclusive evidence that actions on GABA systems are involved in the production of its clinical effects. In studies with animals baclofen has been shown to have general CNS depressant properties as indicated by the production of sedation with tolerance, somnolence, ataxia, and respiratory and cardiovascular depression. Baclofen is rapidly and extensively absorbed and eliminated. Absorption may be dose-dependent, being reduced with increasing doses. Baclofen is excreted primarily by the kidney in unchanged form and there is relatively large intersubject variation in absorption and/or elimination.

INDICATIONS AND USAGE

Baclofen tablets, USP are useful for the alleviation of signs and symptoms of spasticity resulting from multiple sclerosis, particularly for the relief of flexor spasms and concomitant pain, clonus, and muscular rigidity.

Patients should have reversible spasticity so that baclofen treatment will aid in restoring residual function. Baclofen tablets, USP may also be of some value in patients with spinal cord injuries and other spinal cord diseases.

Baclofen tablets, USP are not indicated in the treatment of skeletal muscle spasm resulting from rheumatic disorders.

The efficacy of baclofen in stroke, cerebral palsy, and Parkinson's disease has not been established and, therefore, it is not recommended for these conditions.

CONTRAINDICATIONS

Hypersensitivity to baclofen.

WARNINGS

a. **Abrupt Drug Withdrawal:** Hallucinations and seizures have occurred on abrupt withdrawal of baclofen. Therefore, except for serious adverse reactions, the dose should be reduced slowly when the drug is discontinued.

b. **Impaired Renal Function:** Because baclofen is primarily excreted unchanged through the kidneys, it should be given with caution, and it may be necessary to reduce the dosage.

c. **Stroke:** Baclofen has not significantly benefited patients with stroke. These patients have also shown poor tolerability to the drug.

d. **Pregnancy:** Baclofen has been shown to increase the incidence of omphaloceles (ventral hernias) in fetuses of rats given approximately 13 times the maximum dose recommended for human use, at a dose which caused significant reductions in food intake and weight gain in dams. This abnormality was not seen in mice or rabbits.

There was also an increased incidence of incomplete sternebral ossification in fetuses of rats given approximately 13 times the maximum recommended human dose, and an increased incidence of unossified phalangeal nuclei of forelimbs and hindlimbs in fetuses of rabbits given approximately 7 times the maximum recommended human dose. In mice, no teratogenic effects were observed, although reductions in mean fetal weight with consequent delays in skeletal ossification were present when dams were given 17 and 34 times the human daily dose. There are no studies in pregnant women. Baclofen should be used during pregnancy only if the benefit clearly justifies the potential risk to the fetus.

PRECAUTIONS

Because of the possibility of sedation, patients should be cautioned regarding the operation of automobiles or other dangerous machinery, and activities made hazardous by decreased alertness. Patients should also be cautioned that the central nervous system effects of baclofen may be additive to those of alcohol and other CNS depressants.

Baclofen should be used with caution where spasticity is utilized to sustain upright posture and balance in locomotion or whenever spasticity is utilized to obtain increased function.

In patients with epilepsy, the clinical state and electroencephalogram should be monitored at regular intervals, since deterioration in seizure control and EEG have been reported occasionally in patients taking baclofen.

It is not known whether this drug is excreted in human milk. As a general rule, nursing should not be

undertaken while a patient is on a drug since many drugs are excreted in human milk.

A dose-related increase in incidence of ovarian cysts and a less marked increase in enlarged and/or hemorrhagic adrenal glands was observed in female rats treated chronically with baclofen.

Ovarian cysts have been found by palpation in about 4% of the multiple sclerosis patients that were treated with baclofen for up to one year. In most cases these cysts disappeared spontaneously while patients continued to receive the drug. Ovarian cysts are estimated to occur spontaneously in approximately 1% to 5% of the normal female population.

Pediatric Use

Safety and effectiveness in pediatric patients below the age of 12 years have not been established.

ADVERSE REACTIONS

The most common is transient drowsiness (10 to 63%). In one controlled study of 175 patients, transient drowsiness was observed in 63% of those receiving baclofen compared to 36% of those in the placebo group. Other common adverse reactions are dizziness (5 to 15%), weakness (5 to 15%) and fatigue (2 to 4%).

Others reported:

Neuropsychiatric: Confusion (1 to 11%), headache (4 to 8%), insomnia (2 to 7%); and, rarely, euphoria, excitement, depression, hallucinations, paresthesia, muscle pain, tinnitus, slurred speech, coordination disorder, tremor, rigidity, dystonia, ataxia, blurred vision, nystagmus, strabismus, miosis, mydriasis, diplopia, dysarthria, epileptic seizure.

Cardiovascular: Hypotension (0 to 9%). Rare instances of dyspnea, palpitation, chest pain, syncope.

Gastrointestinal: Nausea (4 to 12%), constipation (2 to 6%); and rarely, dry mouth, anorexia, taste disorder, abdominal pain, vomiting, diarrhea, and positive test for occult blood in stool.

Genitourinary: Urinary frequency (2 to 6%); and rarely, enuresis, urinary retention, dysuria, impotence, inability to ejaculate, nocturia, hematuria.

Other: Instances of rash, pruritus, ankle edema, excessive perspiration, weight gain, nasal congestion.

Some of the CNS and genitourinary symptoms may be related to the underlying disease rather than to drug therapy. The following laboratory tests have been found to be abnormal in a few patients receiving baclofen: increased SGOT, elevated alkaline phosphatase, and elevation of blood sugar.

OVERDOSAGE

Signs and Symptoms: Vomiting, muscular hypotonia, drowsiness, accommodation disorders, coma, respiratory depression and seizures.

Treatment: In the alert patient, empty the stomach promptly by induced emesis followed by lavage. In the obtunded patient, secure the airway with a cuffed endotracheal tube before beginning lavage (do not induce emesis). Maintain adequate respiratory exchange, do not use respiratory stimulants.

DOSAGE AND ADMINISTRATION

The determination of optimal dosage requires individual titration. Start therapy at a low dosage and increase gradually until optimum effect is achieved (usually between 40-80 mg daily).

The following dosage titration schedule is suggested:

5 mg t.i.d. for 3 days

10 mg t.i.d. for 3 days

15 mg t.i.d. for 3 days

20 mg t.i.d. for 3 days

Thereafter additional increases may be necessary but the total daily dose should not exceed a maximum of 80 mg daily (20 mg q.i.d.).

The lowest dose compatible with an optimal response is recommended. If benefits are not evident after a reasonable trial period, patients should be slowly withdrawn from the drug (See **WARNINGS**, Abrupt Drug Withdrawal).

HOW SUPPLIED

Baclofen Tablets, USP 5 mg are available as a White to off white, round flat-faced, beveled edge tablets debossed with "023" on one side and plain on other side, containing 5 mg baclofen USP packaged in bottles of 100, 500 and 1000 tablets.

NDC 52817-319-10 100s count

Baclofen Tablets, USP 10 mg are available as a White to off white, round flat-faced, beveled edge tablets debossed with "024" on one side and score on other side, containing 10 mg baclofen USP packaged in bottles of 100, 500 and 1000 tablets.

NDC 52817-320-10 100s count

NDC 52817-320-50 500s count

NDC 52817-320-00 1000s count

Baclofen Tablets, USP 20 mg are available as a White to off white, round flat-faced, beveled edge tablets debossed with "025" on one side and score on other side, containing 20 mg baclofen USP packaged in bottles of 100, 500 and 1000 tablets.

NDC 52817-321-10 100s count

NDC 52817-321-50 500s count

NDC 52817-321-00 1000s count

PHARMACIST: Dispense in a well closed container as defined in the USP, with a child-resistant closure (as required).

Store at 20° to 25°C (68°to 77°F) [See USP Controlled Room Temperature].

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Distributed by:

TruPharma, LLC

Tampa, FL 33609

Manufactured by:

Rubicon Research Private Limited,

Ambernath, Dist: Thane, 421506

India

Rev.: 01, February 2018

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC (52817-319-10)

100 Tablets

Baclofen Tablets, USP 5 mg

Rx only

Rubicon Research Private Limited

Each tablet contains: Baclofen, USP 5 mg	Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].	NDC 52817- 319 -10	Distributed by: TruPharma, LLC Tampa, FL 33609	 N 31 52817 31910 14	
Usual Dosage: See package insert for full prescribing information.	KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN	Baclofen Tablets, USP 5 mg	Manufactured by: Rubicon Research Private Limited Ambernath, Dist. Thane, 421506 India		
Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).	Rev.02, 05/18	Rx Only 100 Tablets	Unvarnished Zone 45 x 15 mm		
					

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL




NDC (52817-320-10)

100 Tablets

Baclofen Tablets, USP 10 mg

Rx only

Rubicon Research Private Limited

Each tablet contains: Baclofen, USP 10 mg	Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].	NDC 52817- 320 -10	Distributed by: TruPharma, LLC Tampa, FL 33609	 N 31 52817 32010 10	
Usual Dosage: See package insert for full prescribing information.	KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN	Baclofen Tablets, USP 10 mg	Manufactured by: Rubicon Research Private Limited Ambernath, Dist. Thane, 421506 India		
Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).	Rev.02, 05/18	Rx Only 100 Tablets	Unvarnished Zone 45 x 15 mm		
					

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL




NDC (52817-320-50)

500 Tablets

Baclofen Tablets, USP 10 mg

Rx only

Rubicon Research Private Limited

<p>Each tablet contains: Baclofen, USP 10 mg</p> <p>Usual Dosage: See package insert for full prescribing information. Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required). Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].</p> <p>KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN</p> <p>Rev.02, 04/19</p> 	<p>NDC 52817-320-50</p> <div><p>Baclofen Tablets, USP 10 mg</p></div> <p>Rx Only 500 Tablets</p> 	<p>Distributed by: Trupharma, LLC Tampa, FL 33609</p> <p>Manufactured by: Rubicon Research Private Limited Ambernath, Dist. Thane, 421506 India Mfg Lic No. : KD-682 PMS0588</p>  <p>N 3 52817 32050 6</p>
<p>Unvarnished Zone 50 x 17 mm</p>		

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL



NDC (52817-320-00)

1000 Tablets

Baclofen Tablets, USP 10 mg

Rx only

Rubicon Research Private Limited

Each tablet contains: Baclofen, USP 10 mg	NDC 52817- 320 -00	Distributed by: TruPharma, LLC Tampa, FL 33609 Manufactured by: Rubicon Research Private Limited Ambarnath, Dist. Thane, 421506 India Mfg Lic No. : KD-682 PMS0512	
Usual Dosage: See package insert for full prescribing information. Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required). Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN Rev.02, 05/18	<div data-bbox="639 153 964 436" style="background-color: #007bff; color: white; padding: 10px; text-align: center;"> Baclofen Tablets, USP 10 mg </div> <div data-bbox="639 449 964 480" style="display: flex; justify-content: space-between;"> Rx Only 1000 Tablets </div> <div data-bbox="680 600 924 651" style="text-align: center;">  </div>		Unvarnished Zone 48 x 17 mm

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL


NDC (52817-321-10)

100 Tablets

Baclofen Tablets, USP 20 mg

Rx only

Rubicon Research Private Limited

Each tablet contains: Baclofen, USP 20 mg	NDC 52817- 321 -10	Distributed by: TruPharma, LLC Tampa, FL 33609 Manufactured by: Rubicon Research Private Limited Ambarnath, Dist. Thane, 421506 India	
Usual Dosage: See package insert for full prescribing information. Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required). Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN	<div data-bbox="501 1249 773 1486" style="background-color: #ff7f0e; color: white; padding: 10px; text-align: center;"> Baclofen Tablets, USP 20 mg </div> <div data-bbox="501 1514 773 1543" style="display: flex; justify-content: space-between;"> Rx Only 100 Tablets </div> <div data-bbox="526 1675 748 1724" style="text-align: center;">  </div>		Mfg Lic No. : KD-682 PMS0511 Rev.02, 05/18

Unvarnished Zone
52 x 22 mm

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL



NDC (52817-321-50)

500 Tablets

Baclofen Tablets, USP 20 mg

Rx only

Rubicon Research Private Limited

Each tablet contains: Baclofen, USP 20 mg	NDC 52817- 321 -50	Distributed by: Trupharma, LLC Tampa, FL 33609	 3 52817 32150 3
Usual Dosage: See package insert for full prescribing information. Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required). Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].	Baclofen Tablets, USP 20 mg	Manufactured by: Rubicon Research Private Limited Ambernath, Dist. Thane, 421506 India Mfg Lic No. : KD-682 PMS0589	
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN Rev.02, 04/19	Rx Only 500 Tablets	Unvarnished Zone 50 x 17 mm	
			

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL




NDC (52817-321-00)

1000 Tablets

Baclofen Tablets, USP 20 mg

Rx only

Rubicon Research Private Limited

Each tablet contains: Baclofen, USP 20 mg	NDC 52817- 321 -00	Distributed by: TruPharma, LLC Tampa, FL 33609	 3 52817 32100 8
Usual Dosage: See package insert for full prescribing information. Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required). Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].	Baclofen Tablets, USP 20 mg	Manufactured by: Rubicon Research Private Limited Ambernath, Dist. Thane, 421506 India Mfg Lic No. : KD-682 PMS0513	
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN Rev.02, 05/18	Rx Only 1000 Tablets	Unvarnished Zone 60 x 20 mm	
			

BACLOFEN

baclofen tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:528 17-319
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)	BACLOFEN	5 mg

Inactive Ingredients	
Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics			
Color	white (White to off white)	Score	no score
Shape	ROUND	Size	5mm
Flavor		Imprint Code	023
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:528 17-319-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209102	04/01/2018	

BACLOFEN			
baclofen tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:528 17-320
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)	BACLOFEN	10 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	white (White to off white)	Score	2 pieces
Shape	ROUND	Size	7mm
Flavor		Imprint Code	024
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52817-320-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2018	
2	NDC:52817-320-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2018	
3	NDC:52817-320-00	1000 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209102	04/01/2018	

BACLOFEN

baclofen tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:52817-321
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)	BACLOFEN	20 mg

Inactive Ingredients				
Ingredient Name			Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
STARCH, CORN (UNII: O8232NY3SJ)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
Product Characteristics				
Color	white (White to off white)	Score	2 pieces	
Shape	ROUND	Size	9mm	
Flavor		Imprint Code	025	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52817-321-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2018	
2	NDC:52817-321-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2018	
3	NDC:52817-321-00	1000 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2018	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA209102		04/01/2018	

Labeler - TruPharma LLC (078533947)

Registrant - Rubicon Research Private Limited (918629544)

Establishment			
Name	Address	ID/FEI	Business Operations
Rubicon Research Private Limited		677604197	manufacture(52817-319, 52817-320, 52817-321) , analysis(52817-319, 52817-320, 52817-321)

Revised: 1/2021

TruPharma LLC